

Lecture Presentation

Chapter 6

Ionic Compounds: Periodic Trends & Bonding Theory

HW: 6.1, 6.2, 6.3, 6.5, 6.6,
6.7, 6.9, 6.10, 6.11, 6.12,
6.13, 6.16, 6.20, 6.22, 6.24,
6.26, 6.38, 6.40, 6.46, 6.52,
6.62, 6.68

John E. McMurry
Robert C. Fay

Electron Configurations of Ions (for main group elements)

stable ion = closest noble gas electron configuration
(to octet)



metal – lose electrons (group 1A to group 3A)
(**+1** lose one electron to **+3** lose 3 electrons)

non metal – gain electrons (group 5A to group 7A)
(**-3** gain 3 electrons to **-1** gain 1 electron)


+ charge = remove electron until get electron
configuration of closest prior noble gas

- charge = add electron until get electron
configuration of closest next noble gas

Electron Configurations of Ions



Electron Configurations of Ions



Electron Configurations of Ions

TABLE 6.1 Some Common Main-Group Ions and Their Noble-Gas Electron Configurations

Group 1A	Group 2A	Group 3A	Group 6A	Group 7A	Electron Configuration
H ⁺					[None]
H ⁻					[He]
Li ⁺	Be ²⁺				[He]
Na ⁺	Mg ²⁺	Al ³⁺	O ²⁻	F ⁻	[Ne]
K ⁺	Ca ²⁺	*Ga ³⁺	S ²⁻	Cl ⁻	[Ar]
Rb ⁺	Sr ²⁺	*In ³⁺	Se ²⁻	Br ⁻	[Kr]
Cs ⁺	Ba ²⁺	*Tl ³⁺	Te ²⁻	I ⁻	[Xe]

*These ions don't have a true noble-gas electron configuration because they have an additional filled *d* subshell.

HW: Electron Configurations of Ions (for main group elements)

metal – lose electrons to nearest noble gas (+ charge)

non metal – gain electrons to nearest noble gas (- charge)

What is the electron configuration of F ?

What is the electron configuration of F^{-1} ?

What is the electron configuration of Mg ?


What is the electron configuration of Mg^{+2} ?

Electron Configurations of Ions (transition metals)

stable ion = lose electrons to half filled d or lose only s electrons (for early TM), usually lose s electron before d electron

Many transition metals have multiple charges possible

Electron Configurations of Ions



HW: Electron Configurations of Ions (for transition metal elements)

Usually lose s electrons first. Lose d electrons to half full or empty d subshell.

What is the electron configuration of Mn ?


What is the electron configuration of Mn^{+2} ?

What is the electron configuration of Zn ?


What is the electron configuration of Zn^{+2} ?

Ionic Radii

(a)



(b)




Cations are smaller than the corresponding neutral atoms, both because the principal quantum number of the valence-shell electrons is smaller for the cations than it is for the neutral atoms and because Z_{eff} is larger.

cations are smaller than neutral atoms (keep same nucleus but fewer electrons)

because fewer electrons to lower n (principal quantum number) & larger Z_{eff}

End 10/28 Monday F section

Ionic Radii



Anions are larger than their **neutral atoms** because of additional electron–electron repulsions and a decrease in Z_{eff} .

Anions – keep same nucleus but larger number of electrons

HW: Ionic Radii

Cations (+ charged) are smaller than neutral atoms

Anions (- charged) are larger than neutral atoms

Which is larger ?

Ca vs Ca^{+2}

S vs S^{-2}

Fe vs Fe^{+2} vs Fe^{+3}

HW: Ionic Radii

Cations (+ charged) are smaller than neutral atoms

Anions (- charged) are larger than neutral atoms

Which is larger ?

Ca vs Ca^{+2} Ca


S vs S^{-2} S^{-2}

Fe vs Fe^{+2} vs Fe^{+3} $\text{Fe} > \text{Fe}^{+2} > \text{Fe}^{+3}$


Ionization Energy



Ionization Energy (E_i): The amount of energy necessary to remove the highest-energy electron from an isolated neutral atom in the gaseous state



Ionization Energy – largest (E_i) is noble gases




Ionization Energy

bc of stability of $\frac{1}{2}$ p subshell

bc of stability of full s subshell

The **group 2A elements (Be, Mg, Ca)** have slightly larger E_i values than might be expected.

The **group 6A elements (O, S)** have slightly smaller E_i values than might be expected.



Boron has a lower E_i due to a smaller Z_{eff} (shielding by the 2s electrons or remove only e in p subshell).

Higher Ionization Energies (zig zag line - big jump in ionization energy if have to remove core electrons – to lower shell)

End 10/28 M G section



First ionization E_{i1}



second ionization E_{i2}



Third ionization E_{i3}

TABLE 6.2 Higher Ionization Energies (kJ/mol) for Main-Group Third-Row Elements

Group	1A	2A	3A	4A	5A	6A	7A	8A
E_i Number	Na	Mg	Al	Si	P	S	Cl	Ar
E_{i1}	496	738	578	787	1,012	1,000	1,251	1,520
E_{i2}	4,562	1,451	1,817	1,577	1,903	2,251	2,297	2,665
E_{i3}	6,912	7,733	2,745	3,231	2,912	3,361	3,822	3,931
E_{i4}	9,543	10,540	11,575	4,356	4,956	4,564	5,158	5,770
E_{i5}	13,353	13,630	14,830	16,091	6,273	7,013	6,540	7,238
E_{i6}	16,610	17,995	18,376	19,784	22,233	8,495	9,458	8,781
E_{i7}	20,114	21,703	23,293	23,783	25,397	27,106	11,020	11,995

The zigzag line marks the large jumps in ionization energies.